

Effect of supplementation fat during the last 3 weeks of uterine life and the preweaning period on performance, ruminal fermentation, blood metabolites, passive immunity and health of the newborn calf

A. R. Jolazadeh¹, T. Mohammadabadi^{1*}, M. Dehghan-banadaky², M. Chaji¹ and M. Garcia^{3†}

¹Department of Animal Science, Faculty of Animal Science and Food Technology, Agricultural Sciences and Natural Resources University of Khuzestan, P.O. Box 63517-73637, Mollasani, Abvaz, Iran

²Department of Animal Science, Campus of Agriculture and Natural Resources, University of Tebran, Karaj, Iran

³Department of Animal Science and Industry, Kansas State University, Manhattan, KS 66502, USA

(Submitted 8 March 2019 – Final revision received 28 July 2019 – Accepted 16 August 2019)

Abstract

The objective of this study was to evaluate the effects of supplementing essential fatty acids (FA), during late gestation and the preweaning and early weaning periods on passive immunity, growth, health, rumen fermentation parameters, blood metabolites and behaviour of dairy calves. During the last 3 weeks of pregnancy, cattle (*n* 120), within parity, were randomly assigned to one of three diets with different fat supplements: (a) no supplemental fat (CON), (b) supplement rich in linoleic acid (CSO), or (c) supplement rich in EPA and DHA (CFO). Eighty-four newborn Holstein calves were randomly assigned, within the prepartum diets, to one of two calf starters: no fat supplement (FC-0) or 2% Ca-salt of unsaturated FA (FC-2). Overall, the interaction between dam diets and calf starters did not affect calf performance or any other parameter measured. Calves born from dams fed fat (CSO or CFO) performed better than calves born from dams fed CON. Namely, calves born from dams fed fat had greater plasma concentrations of IgG ($P < 0.01$), better apparent efficiency of IgG absorption ($P < 0.01$) and average daily gain (ADG, 597 *v.* 558 g/d; $P = 0.02$), and lower rectal temperature (RT; $P < 0.01$). Calves fed a calf starter rich in unsaturated FA (FC-2) had greater ($P \leq 0.01$) ADG, skeletal growth, feed efficiency, and weaning weight compared with FC-0-fed calves. Furthermore, calves fed FC-2 had lower RT during the pre- and post-weaning periods ($P \leq 0.04$) and fewer days with diarrhoea ($P < 0.001$) compared with calves fed FC-0. Time spent eating, ruminating, standing, lying, and on non-nutritive oral behaviour did not differ by treatment. Similarly, treatments did not affect ruminal fermentation parameters. At 28 and 77 d of age, calves fed FC-2 had higher plasma concentrations of albumin and cholesterol ($P \leq 0.02$) and lower urea N compared with calves fed FC-0. Plasma concentrations of alkaline phosphatase were higher in calves fed FC-2 compared with those fed FC-0, when they were 77 d old. **These findings support feeding moderate amounts of long-chain PUFA during late uterine life or during the preweaning period have beneficial effects on calf metabolism, growth, and health performance.**

Key words: Holstein calves: Performance: Fatty acids: Peripartum diet

The cost of rearing replacement heifers is one of the highest investments in the dairy industry. Thus, proper nutritional and management practices must be placed to maximise the return of investment. Particularly, during the preweaning and early weaning period, the most critical management goals are to minimise the incidence of diseases and mortality and maximise growth. Several studies have concluded that the benefits of proper preweaning management, leading to maximise average

daily gain (ADG), have not only short but also lifelong impacts on heifers' performance^(1,2).

The discovery of the essentiality of fatty acids (FA), namely linoleic⁽³⁾ and α -linolenic acids⁽⁴⁻⁶⁾, led the livestock scientific community to further investigate the role of these essential FA (EFA) and their derivatives arachidonic acid (C:20 : 4*n*-6) and EPA (20 : 5*n*-3) and DHA (22 : 6*n*-3), respectively. FA are more than structural components of TAG; as components of cell

Abbreviations: ADG, average daily gain; AEA, apparent efficiency of IgG absorption; ALP, alkaline phosphatase; BW, body weight; CFO, Ca-salts of fish oil; CON, control (no supplemental fat); CP, crude protein; CS, calf starter; CSO, Ca-salts of soyabean oil; DD, dam diet; DMI, DM intake; EFA, essential fatty acid; FA, fatty acid; FAT, contrast of fat supplement *v.* no fat supplement (CSO + CFO *v.* CON); FC, fat Ca-salts; FC-0, calf starter with no fat supplement; FC-2, calf starter supplemented with 2% fat Ca-salt; FE, feed efficiency; MR, milk replacer; RT, rectal temperature; SFO, contrast of fatty acid source (CSO *v.* CFO); VFA, volatile fatty acid.

* **Corresponding author:** T. Mohammadabadi, email mohammadabadi@asnrkh.ac.ir; t.mohammadabadi.t@gmail.com

† Present address: Phibro Animal Health Corporation, Teaneck, NJ 07666, USA.

membranes, they regulate signal transduction⁽⁷⁾. Several studies in humans and rodents have reported that *n*-3 compared with *n*-6 FA are more potent activators of genes involved in lipid oxidation, more potent reducers of genes encoding enzymes for lipid synthesis and important regulators of immune response^(8,9).

Studies have confirmed that the strategic feeding of FA enriches tissues and body fluids with the corresponding supplemented FA. In dairy cattle, dams supplemented with prepartum diets enriched in linoleic acid⁽¹⁰⁾ or EPA and DHA⁽¹¹⁾ produced colostrum with greater concentrations of the respective supplemented FA compared with colostrum from non-supplemented cows. Furthermore, plasma of newborn dairy calves born from dams fed diets enriched in EPA and DHA⁽¹²⁾ or linoleic acid⁽¹³⁾ had increased concentrations of the FA fed to their dams. In beef cattle, plasma (30 or 60 d of age) and adipose tissue (60 d of age) of calves suckling dams supplemented with high-oleic or high-linoleic safflower seeds had increased concentrations of the supplemented FA in plasma and adipose tissue, compared with calves suckling non-supplemented dams⁽¹⁴⁾. Furthermore, pre-weaned dairy calves fed milk replacer (MR) supplemented with EFA also reported increased concentrations of the corresponding EFA and/or that of their derivatives in plasma^(13,15,16) and liver^(16,17).

Previous studies have reported the benefits of feeding diets enriched with EFA during late uterine life on performance of newborn calves. Salehi *et al.*⁽¹⁸⁾ reported better concentrations of colostrum IgG and Garcia *et al.*⁽¹³⁾ reported better efficiency of IgG absorption of calves born from dams fed diets enriched with EFA. Adequate passive transfer of IgG is crucial to minimise neonatal morbidity and mortality and strengthen calf immunity⁽¹⁹⁾. Calves born from dams supplemented with long-chain FA during the prepartum may have enterocytes with a particular FA profile that could benefit passive transfer of immunoglobulins⁽¹⁰⁾. Furthermore, strategic supplementation of EFA in MR^(13,20) or calf starter (CS)^(21,22) fed to preweaned calves has been reported to impact calf metabolism, improve growth, feed efficiency (FE), health, and immune status.

The hypothesis of this study was that supplementing prepartum diets with Ca-salts of soyabean oil (CSO) or Ca-salts of fish oil (CFO) would improve immune status, health and growth of newborn calves compared with calves born from dams non-fat-supplemented. Another hypothesis was that calves born from dams supplemented with EFA or their derivatives would benefit to a greater extent from eating a CS supplemented with Ca salts of unsaturated FA. Therefore, the objective was to evaluate the effect of supplementing diets enriched in linoleic acid or EPA and DHA to late-gestation Holstein cattle on metabolic profile, immune status, health and growth of calves consuming starter diets enriched in oleic and linoleic acids during pre- and postweaning periods.

Methods

Dam management and dietary treatments

The experiment was conducted from July 2016 through October 2016, at a commercial dairy farm (Zarin Khoshe Dairy Farm). All

experimental procedures were conducted according to The Care and Use of Agricultural Animals in Research and Teaching⁽²³⁾ guidelines. The Animal Experiment Committee at Tehran University, Karaj, Iran, approved all procedures and guidelines involving animals. The present study was part of a large research project, with a portion of the results (prepartum and postpartum performance of calves' dams) reported elsewhere⁽¹¹⁾. Briefly, pregnant nulliparous (*n* 42) and previously parous (*n* 78) Holstein cattle were blocked by parity and enrolled in the study starting at 3 week before their calculated parturition date. The basal prepartum diet was formulated to have low concentrations of total FA (1.87% of DM), linoleic acid (0.84% of DM) and α -linolenic acid (0.36% of DM). Prepartum cattle were fed one of the following three diets: (1) no fat supplement (CON), (2) 1.15% of dietary DM as CSO (140 g/cattle per d) supplement (Persiafat; Kimiya Danesh Alvand Co.) or (3) 1.15% of dietary DM as CFO (140 g/cattle per d) supplement (Persiafat; Kimiya Danesh Alvand Co.). The CSO and SFO diets were isoenergetic but had different FA profiles (linoleic acid: 1.30 and 0.87%; EPA and DHA: 0 and 0.15% for CSO and SFO, respectively). All diets were isonitrogenous. Ingredient and chemical composition of the prepartum diets and specific details on prepartum feeding and management were reported previously⁽¹¹⁾.

Calving management and colostrum, milk and starter feeding

Calves were born in a sod-based pen. All dams were monitored for signs of calving initiation every 30 min between 05.30 and 15.30 hours and then every 2 h between 15.30 and 05.30 hours. Within 2 h of birth, calves were weighed and ear-tagged, and the umbilical cord was disinfected with 10% Betadine solution (Purdue Frederick Co.). Calves were housed temporarily in individual pens (1 × 1 m) equipped with a heat lamp before moving to individual wired pens (1 × 1.5 m) bedded with sand between 6 and 16 h of age. Cows were milked with a cow-side vacuum pump within 1 h after calving. Colostrum from each cow (4 litres) was fed to their corresponding calf, regardless of IgG concentration. Remnant colostrum (>1 litre and >50 g of IgG/l) was stored at -20°C to feed calves born from dams fed the same dietary treatment but producing low amounts of colostrum. Calves were considered to have adequate passive transfer if the plasma concentration of total IgG was ≥ 10 g/l after 24 h of colostrum feeding⁽²⁴⁾. Whole colostrum samples were collected and frozen (-20°C) for later analyses of IgG concentration and FA profile, and results were reported elsewhere⁽¹¹⁾. Briefly, the FA profile of colostrum resembled that of the prepartum diets fed to dams during late gestation⁽¹¹⁾. Furthermore, given that an extensive body of research has confirmed that the FA profile of tissues and body fluids resembles that of the supplemented diets, coupled with budgetary restrictions, we did not measure the FA profile of tissues or body fluids of calves in the present study. Plasma samples of newborn calves (at birth and 24 h after colostrum feeding) were measured for bovine total IgG concentration using radioimmunoassay⁽¹⁰⁾. Intake of IgG was calculated using the concentration of colostrum IgG reported previously⁽¹¹⁾. The apparent efficiency of IgG absorption (AEA, %) was calculated according to Quigley *et al.*⁽²⁵⁾ using the following



Table 1. Fatty acid (FA) profile (g/100 g of total FA) of calf starters and fat supplement used

FA	Fat supplement	Calf starters	
	FC*	FC-0	FC-2
C16 : 0	28	14.5	19.2
C16 : 1	3.0	0.09	1.10
C18 : 0	5.0	2.54	3.37
C18 : 1	26	21.57	23.63
C18 : 2 <i>n</i> -6	30	55.30	46.60
C18 : 3 <i>n</i> -3	3.0	3.80	3.60
Other	5.0	2.20	2.50
Σ SFA	33	17.04	22.57
Σ UFA	62	80.76	74.93
Σ PUFA	33	59.1	50.2

FC, fat Ca-salts; FC-0, calf starter with no fat supplement; FC-2, calf starter supplemented with 2 % fat Ca-salt; UFA, unsaturated FA.

* Fat Ca-salt rich in unsaturated FA (Persiafat, Kimiya Danesh Alvand Co.).

equation: IgG concentration in plasma at 24 h of life (g/l) × (0.099 × body weight (BW) (kg) at birth)/IgG intake (g). Plasma volume was estimated at 9.9 % of birth weight⁽²⁵⁾. Eighty-four Holstein calves were used in the present study (*n* 14 calves per factorial treatment, ten males and four females) in a completely randomised design, with dietary treatments in a 3 × 2 factorial arrangement (three dam diets (DD) and two CS; *n* 6). Calves were fed warm (38°C) pasteurised tank milk twice per d (08.00 and 17.00 hours) using steel buckets, starting at 3 d of age, and always drank all their offered milk. Volumes of milk fed decreased with age: 4 litres from day 4 to 7, and then 5, 6, 7, 6.5, 5.5, 4 and 2 litres during weeks 2 to 8, respectively. The average nutritional composition of milk was 11.50 % DM, 2.98 % crude protein (CP), 3.35 % fat and 4.51 % lactose. Calves had *ad libitum* access to one of two CS: (1) no fat supplement (FC-0) or (2) supplemented with 2 % fat Ca-salt rich in unsaturated FA, primarily oleic and linoleic acids (FC-2; Persiafat, Kimiya Danesh Alvand Co.; Table 1). All calves were weaned at 56 d but the experimental CS were fed until 77 d of age. CS were isonitrogenous, and the ingredients and composition are shown in Table 2. All calves had free access to clean and freshwater at all times. The offered starter was adjusted daily to achieve 5–10 % refusals (i.e. the portion of the starter not consumed over a 24-h period); and refusals were collected and weighed daily at 08.00 hours. Chopped high-quality alfalfa hay (5 % of CS) was included in the starter diets once calves were 30 d of age. The starter diets (CS or CS + alfalfa hay) were offered once daily at 09.00 hours.

Data collection and sampling

All calves (*n* 84) were weighed at birth, 56 d and 77 d of age to measure changes in BW and calculate ADG. In addition, all calves were weighed once a week to observe weekly growth patterns. Body measurements of all calves were taken at 3 d of age, weaning (d 56) and the end of the study (d 77), in the morning and before feeding according to Lesmeister & Heinrichs⁽²⁶⁾. Body measurements included heart girth (circumference of the chest), hip width (distance between the points of hook bones), body length (distance between the points of

Table 2. Ingredients and nutrient composition (g/kg DM unless otherwise noted) of calf starters

Item	Calf starter	
	FC-0	FC-2
Ingredient (g/kg DM)		
Maize, ground	500	480
Barley, ground	100	100
Maize gluten meal	40	40
Beet pulp dry	90	90
Extruded soyabean seed	20	20
Soyabean meal	200	200
Fat Ca-salts*	0	20
Calcium carbonate	7.0	7.0
Sodium bicarbonate	10	10
Vitamin–mineral premix†	10	10
White salt	5.0	5.0
Monensin sodium 10 %	1.0	1
Sodium bentonite	12	12
Dicalcium phosphate	5.0	5.0
Nutrient composition (g/kg DM)		
CP	20.10	19.92
NDF	14.85	14.62
ADF	7.82	7.71
NFC‡	53.80	52.40
Ether extract	3.35	5.10
ME§ (MJ/kg DM)	11.13	11.55
NEM§ (MJ/kg DM)	7.24	7.65
NEg§ (MJ/kg DM)	4.64	5.06

FC-0, calf starter with no fat supplement; FC-2, calf starter supplemented with 2 % fat Ca-salt; CP, crude protein; NDF, neutral-detergent fibre; ADF, acid-detergent fibre; NFC, non-fibre carbohydrates; ME, metabolisable energy, NEM, net energy for maintenance; NEg, net energy for gain.

* Fat Ca-salts, Persiafat, Kimiya Danesh Alvand Co., contained 85 % fat and 9 % Ca (1 % C14:0, 28 % C16 : 0; 3 % C16 : 1; 5 % C18 : 0, 26 % C18 : 1, 30 % C18 : 2; 3 % C18 : 3, 4 % other).

† Vitamin–mineral premix contained per kg: Ca 160 g; Mg 40 g; Na 30 g; Fe 3 g; Cu 3 g; Zn 7 g; Mn 4 g; I 0.08 g; Co 0.05 g; Se 0.06 g; vitamin A 800 000 IU; vitamin D 150 000 IU; vitamin E 2000 IU.

‡ Calculated as DM – (NDF + CP + ether extract + ash) (National Research Council, 2001).

§ Estimated using National Research Council (2001) equations.

shoulder and rump), body barrel (circumference of the belly before feeding), wither height (distance from the base of the front feet to wither) and hip height (distance from the base of the rear feet to the hook bones).

Amount of starter diets offered and refused per calf were recorded daily throughout the experiment. Feed samples were collected weekly and stored at –20°C for chemical analyses. Subsamples of feeds and refusals were mixed thoroughly, dried at 55°C for 48 h and ground to pass through a 1 mm screen in a hammer mill (Arthur Hill Thomas Co.) before chemical analyses for DM (method 934.01), CP (method 988.05) and ether extract (method 920.39)⁽²⁷⁾. The neutral-detergent fibre contents were analysed according to Van Soest *et al.*⁽²⁸⁾, assayed with sodium sulphite, without α -amylase, and were expressed with residual ash. Eight representative samples of milk were collected throughout the study and analysed for DM, fat, CP and lactose content by infrared spectroscopy (Foss Electric). FE (kg of BW gain/kg of total DM intake (DMI)) was calculated for the preweaning period, days 57–77 and for the extent of the study. Total DMI corresponds to the amount of DMI contributed by the starter diets (CS and alfalfa hay) and by the fixed amounts of milk provided to each calf.

Total tract apparent digestibility of DM, organic matter, CP, neutral-detergent fibre, acid-detergent fibre and ether extract were determined using acid-insoluble ash as an internal marker⁽²⁹⁾. Faecal samples were obtained from each calf twice per d (in the morning and afternoon) during four consecutive days (from days 74 to 77). Faecal samples were dried at 55°C in a forced air oven, ground to pass a 1-mm screen in a Wiley mill (Arthur Hill Thomas Co.) and then mixed thoroughly.

Samples of ruminal fluid (approximately 150 ml) were collected from each calf on days 30 and 70, about 3 h after the morning feeding, using a stomach tube attached to an Erlenmeyer flask and vacuum pump; the first 50 ml were discarded to avoid possible saliva contamination. The ruminal fluid was filtered through four layers of cheesecloth, and pH was determined immediately using a portable pH meter (Sentron, model A102-003). Approximately 25 ml of filtrate were then preserved by adding 5 ml of 25% metaphosphoric acid solution for later determination of volatile FA (VFA), and approximately 20 ml of filtrate were combined with 20 ml 0.2 M HCl for later measurement of NH₃-N concentration. The prepared samples were stored at -20°C and then thawed for analysis of respective compounds. Stored samples were centrifuged at 1200 g for 10 min, and the supernatant fluid was analysed for VFA by GC (model 5890; Hewlett-Packard). Ruminal NH₃-N concentration was determined according to the procedures described by Crooke & Simpson⁽³⁰⁾. Blood samples were collected from each calf immediately after birth, 24 h after colostrum feeding and then 3 h after the morning meal on days 28, 56 and 77 of age. Blood was taken from the jugular vein by evacuated tubes containing K2 EDTA (10.5 mg, Monoject; Sherwood Medical). Samples were immediately chilled and then plasma was obtained by centrifugation at 3000 g for 15 min and stored at -20°C. Plasma concentrations of glucose, total protein, albumin, alkaline phosphatase (ALP), cholesterol and TAG were determined using an automatic microplate Reader (EON-BIOTEK) and commercial kits (Pars Azmoon). Enzymatic assays were used for analyses of plasma insulin (DRG), blood urea N (Pars Azmoon) and NEFA (Randox Laboratories Ltd.) in an automatic spectrophotometer (Clima Plus; RAL) following the manufacturer's instructions. Behavioural data (eating, ruminating, standing, lying and non-nutritive oral behaviour) were monitored by direct observations of each calf in the time (minutes) devoted to each monitored behaviour, once at the last week of the preweaning period (35–42 d) and once at the last week of the postweaning period (70–77 d)⁽³¹⁾. Behavioural data were monitored for 8 consecutive hours following the morning milk feeding (at 08.00 hours) and the CS (09.00 hours) during the last week of the pre- and postweaning periods, respectively.

Calf scoring for health assessment and incidence of health disorders

Attitude, faecal consistency and nasal discharge were scored daily for each calf after the first daily feeding of CS by a single observer using a scale from 0 to 3 (http://www.vetmed.wisc.edu/dms/fapm/fapmtools/8calf/calf_health_scoring_chart.pdf). Attitude scores recorded were (0) responsive, (1) non-active, (2) depressed or (3) moribund. Faecal consistency scores recorded

were (0) firm, no diarrhoea; (1) moderate, soft no diarrhoea; (2) runny faeces, mild diarrhoea; or (3) watery faeces, diarrhoea. Nasal score scale of calves was also scored on a 0–3 scale, with 0 being normal serous discharge, 1 being a small amount of unilateral cloudy discharge, 2 being bilateral cloudy or excessive mucus discharge and 3 being copious bilateral mucopurulent discharge. Calves given a faecal score >1 were considered to have diarrhoea, and those with a score of 3 were considered to have severe diarrhoea. Calves given a score >0 for other criteria were considered as having the disorder. Weekly averages of each score were calculated for statistical analysis. Incidences of health disorders were recorded daily for individual calves. Respiration rate (RR) was measured at 14.00 hours on days 30, 45 and 70 by visual observation for three 1-min periods (not consecutive), according to the methods described by Yari *et al.*⁽³²⁾. Rectal temperature (RT) was measured weekly before morning meals using a rectal thermometer (Yasa Teb Co.), and calves with RT over 39.5°C were categorised as febrile. Calves were vaccinated according to the herd vaccination protocol. Calves with digestive and respiratory problems were treated by farm personnel according to protocols established by the herd veterinarian. All calves remained healthy and exhibited no signs of illness during the experiment.

Statistical analyses

Statistical analyses were conducted for four periods: the first day of life (passive immunity parameters), preweaning (days 0–56), postweaning (days 57–77) and the entire experimental period (days 0–77). Except for starter intake and measures of structural growth, the first measure was performed on day 3 of life. The study was a complete randomised design, with dietary treatments in a factorial arrangement 3 × 2 (3 DD and 2 CS; *n* 6 treatments). The mixed model equation for passive immune variables was defined as $Y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + e_{ijk}$, where Y_{ijk} is the dependent variable, μ is the population mean, α_i is the treatment effect, β_j is the fixed effect of parity, $(\alpha\beta)_{ij}$ is the interaction effect of treatment and parity and e_{ijk} is the residual error. The passive immunity variables were analysed using the PROC MIXED procedure of SAS⁽³³⁾. Contrasts testing the effects of DD were: (1) FAT: control *v.* CFO + SFO and (2) SFO: CSO *v.* CFO, (3) effect of parity (primiparous *v.* multiparous), and the contrasts testing the interactions of DD and parity (4) FAT × parity (contrasts 1 and 3) and (5) SFO × parity (contrasts 2 and 3).

For other continuous variables (rumen fermentation parameters, plasma metabolites, BW, skeletal growth, behaviour data, RT, respiratory rate and apparent nutrient digestibility), the data were analysed using the PROC MIXED procedure of SAS⁽³³⁾. Faecal and health scores were analysed using a multivariable logistic mixed model (GLIMMIX procedure of SAS). The following model was used for variables with repeated-measures, for those with non-repeated-measures the effect of time and its interaction were omitted:

$$Y_{ijklm} = \mu + D_i + C_j + T_k + DC_{ij} + DT_{ik} + CW_{jk} + DCT_{ijk} + \varepsilon_{ijk},$$

where Y_{ijklm} is the observation; μ is overall mean; D_i is the fixed effect of DD (CON, CSO or CFO); C_j is the fixed effect of



Table 3. Passive immunity in calves born from Holstein cows fed prepartum diets non-supplemented (CON) or supplemented with calcium-salts of soyabean oil (CSO) or calcium-salts of fish oil (CFO) fatty acids, during the last 3 weeks before expected parturition (Mean values and pooled standard errors)

Item	Dietary treatment and parity						SEM	<i>P</i> *			
	CON		CSO		CFO			FAT	SFO	Parity	
	Null	Parous	Null	Parous	Null	Parous					
At birth											
TP (g/l)	49.0	43.8	46.5	44.4	47.3	46.0	1.6	0.26	0.45	0.27	
IgG intake† (g)	381	404	481	541	461	532	31.2	<0.01	0.64	0.05	
Plasma IgG (g/l)	0.28	0.26	0.31	0.32	0.30	0.23	0.04	0.64	0.24	0.48	
24 h After birth											
TP (g/l)	53.8	54.4	55.5	56.3	53.8	53.4	1.9	0.71	0.27	0.83	
Plasma IgG (g/l)	18.6	19.9	27.8	28.6	26.2	27.3	1.97	<0.01	0.47	0.50	
Plasma IgG (% of TP)	38.6	40.1	50.8	51.4	49.5	52.2	2.55	<0.01	0.94	0.43	
AEA‡ (%)	22.6	21.8	28.5	26.8	28.6	27.2	1.86	<0.01	0.74	0.32	

CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition; TP, total protein; AEA, apparent efficiency of IgG absorption; BW, body weight.

* FAT = CON *v.* fat supplement; SFO = CSO *v.* CFO. Interactions of FAT and SFO with parity were not significant for any parameter reported in this table ($P > 0.40$).

† IgG intake was calculated using IgG concentration in colostrum, reported in our previous paper⁽¹¹⁾ and the amount offered colostrum was 4 litres, which was similar for all calves.

‡ AEA (%) = (IgG concentration in serum at 24 h of life (g/l) × (0.099 × BW at birth (kg)) / IgG intake (g)) × 100.

CS (FC-0 or FC-2); T_k is the effect of time; DC_{ij} is the interaction of DD × CS; Dt_{jk} is the effect of the interaction between DD and week; Ct_{jk} is the effect of the interaction between CS and week; DCt_{ijk} is the three-way interaction of DD, CS and time; and ϵ_{ijk} is the residual error. Five orthogonal contrasts of DD, CS and interactions were tested for variables measured after initiation of starter feeding. Contrast testing the effects of DD were (1) FAT: control *v.* CFO + SFO, (2) SFO: CSO *v.* CFO, contrast for the effect of CS, (3) FC-0 *v.* FC-2, and the contrast testing the interactions of DD and CS, (4) FAT × CS (contrasts 1 and 3) and (5) SFO × CS (contrasts 2 and 3).

The denominator df were computed using the approximation described by Kenward & Roger⁽³⁴⁾. Significance was declared at $P \leq 0.05$ and trends at $0.05 < P < 0.10$. Tukey's adjustment was used for multiple comparison tests.

Results

Calf absorption of colostral IgG

Total IgG intake was greater in calves born for dams fed fat (CSO or CFO) compared with calves born from dams fed CON (504 *v.* 392 g, $P < 0.01$; Table 3). Concentrations of total IgG in plasma of calves born from dams fed fat supplements compared with calves born from CON-fed dams were 43 and 30 % greater ($P < 0.01$) when expressed as g/l and % total protein, respectively. Calves born from dams fed fat supplements had better ($P < 0.01$) AEA than calves born from CON-fed dams (27.7 *v.* 22.2 %).

Starter intake and growth performance

BW at birth was not impacted by prepartum diets, but BW at weaning (56 d) and 77 d were impacted by prepartum diets (Table 4). Weekly BW, as affected by prepartum diets, started to differentiate at 8 weeks of age, and indeed calves born from dams fed either CSO or CFO were heavier than those born from dams fed CON (Fig. 1(a)). For the preweaning period, calves

born from dams fed fat supplements had greater ADG (542 *v.* 497 g/d; $P = 0.01$; Table 4) compared with calves born from dams fed CON prepartum. Although ADG during the postweaning period was not impacted by prepartum diets, the effect observed during the preweaning period carried on so that the ADG for the total experimental period was also greater for calves born from dams fed fat supplements compared with those born from dams fed CON (597 *v.* 558 g/d; $P = 0.02$). Regardless of DD, calves fed FC-2 had consistently greater ($P < 0.01$) ADG compared with calves fed the FC-0 CS during the pre-weaning (561 *v.* 493 g/d), postweaning (781 *v.* 690 g/d) and overall (621 *v.* 547 g/d) periods. The better ADG of calves fed FC-2 is consistent with their heavier BW, which became significantly different after 6 weeks of age (Fig. 1(b)). Starter intake was not affected by prepartum or starter diets at any evaluated period. FE tended ($P = 0.06$) to be better for preweaned calves born from dams fed fat supplements compared with those born from CON-fed dams (0.55 *v.* 0.51). Regardless of DD, calves fed FC-2 had better ($P < 0.01$) FE compared with calves fed FC-0 CS, during the pre-weaning (0.58 *v.* 0.50), postweaning (0.53 *v.* 0.46) and overall experimental (0.45 *v.* 0.39) periods. Neither DD nor CS or their interactions affected the total tract apparent digestibility of DM, CP, organic matter, EE, acid-detergent fibre and neutral-detergent fibre measured from 74 to 77 d of age.

Data for body measurements are presented in Table 5. Body length, heart girth, hip width, and body barrel were not affected by DD, CS, or their interaction. At 3 d of age, none of the parameters of structural growth was affected by diets or their interaction. Calves born from dams fed fat supplements, compared with those born from dams fed CON, had greater ($P \leq 0.05$) hip (+1 and +0.8 cm, at weaning and the end of the study, respectively) and wither (+0.9 and +1 cm, at weaning and the end of the study, respectively) heights. Regardless of DD, calves fed FC-2, compared with those fed FC-0, had greater ($P < 0.01$) hip (+2.1 and +1.2 cm, at weaning and the end of the study, respectively) and wither (+2.0 and +1.9 cm, at weaning and the end of the study, respectively) heights.

Table 4. Effects of dietary treatments on body weight, starter intake, average daily gain (ADG), gain:feed ratio and apparent digestibility of Holstein calves (*n* 14 per treatment) fed a calf starter (CS) containing no fat supplement (FC-0) or supplemented with 2 % fat calcium-salt (FC-2) in pre- and postweaning periods (Mean values and pooled standard errors)

Items*	Dam diet and calf starters						SEM	P†		
	CON		CSO		CFO			Dam diet		CS
	FC-0	FC-2‡	FC-0	FC-2	FC-0	FC-2		FAT	SFO	
Body weight (kg)										
At birth	40.73	41.20	41.95	39.60	42.13	41.33	0.91	0.72	0.29	0.24
Day 56	67.08	70.52	70.50	72.10	70.14	73.75	1.19	0.03	0.58	<0.01
Day 77	81.40	86.50	85.00	88.61	84.85	90.46	1.33	0.02	0.52	<0.01
ADG (g/d)										
Prewaning	470	524	510	580	500	579	17.3	<0.01	0.74	<0.01
Postweaning	680	760	690	786	700	796	26.5	0.33	0.70	<0.01
Overall	528	588	559	636	555	638	15.2	0.02	0.93	<0.01
Starter intake (g/d)										
Prewaning	413.4	404.5	428.7	407.2	415.8	394.6	23.2	0.67	0.69	0.35
Postweaning	1519	1457	1498	1533	1512	1504	34.0	0.83	0.42	0.68
Overall	754.6	736.6	760.6	755.8	757.6	755.3	29.1	0.84	0.59	0.74
Feed efficiency§										
Prewaning	0.48	0.54	0.51	0.59	0.51	0.60	0.024	0.06	0.97	<0.01
Postweaning	0.45	0.53	0.46	0.53	0.47	0.54	0.021	0.46	0.58	<0.01
Overall	0.38	0.43	0.40	0.46	0.40	0.46	0.016	0.14	0.99	<0.01
Apparent digestibility (g/kg)										
DM (g/kg DM)	816	823	818	823	838	824	10.2	0.29	0.91	0.93
CP	782	783	773	781	779	789	6.90	0.17	0.24	0.14
OM	849	855	863	859	859	861	6.30	0.52	0.16	0.74
NDF	623	617	606	621	631	623	8.50	0.17	0.45	0.92
ADF	515	521	521	531	553	532	13.0	0.14	0.55	0.84
Ether extract	934	933	936	937	934	943	4.80	0.65	0.27	0.16

CON, no fat supplement (control); CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition; CP, crude protein; OM, organic matter; NDF, neutral-detergent fibre; ADF, acid-detergent fibre; DMI, DM intake.

* Starter was offered from 3 d of age. Prewaning: 0–56 d. Postweaning: 57 to 77 d old.

† *P* values for orthogonal contrasts: FAT = control v. CSO + CFO; SFO = CSO v. CFO; CS = FC-0 v. FC-2. Interactions of FAT or SFO with CS were not significant for any parameter reported in this table (*P* ≥ 0.20).

‡ Fat Ca-salts, Persiafat, Kimiya Danesh Alvand Co., contained 85 % fat and 9 % Ca (1 % C14:0, 28 % C16:0; 3 % C16:1; 5 % C18:0, 26 % C18:1, 30 % C18:2; 3 % C18:3, 4 % other).

§ Feed efficiency as calculated from dividing ADG (g) by average daily DMI. DMI includes the DM provided by the milk and starter. The DMI provided by milk was similar among treatments.

Animal behaviour

The total times devoted to different behaviours in an 8-h evaluation period are presented in Table 6. Neither DD nor CS or their interactions affected time spent eating, ruminating, standing, lying or any non-nutritive oral behaviour during the pre- or postweaning periods.

Incidence of diarrhoea and diseases

During the preweaning period, calves born from dams fed fat supplements had a slightly lower (*P* < 0.01) RT compared with those born from CON-fed dams (39.0 v. 39.2°C, respectively; Table 7), but the effect did not extend to the postweaning period. Regardless of DD, calves fed FC-2 had lower (*P* ≤ 0.04) RT compared with calves fed the FC-0 CS during both the pre- (38.97 v. 39.13°C) and postweaning (38.95 v. 39.27°C) periods. Mean scores for attitude and nasal discharge were 0.28 and 0.10 and were not affected by treatments. During the study, calves fed FC-2 had two fewer days with diarrhoea (5.6 v. 7.36 d) compared with those fed FC-0.

Blood metabolites

DD did not affect the mean plasma concentrations of any measured metabolite (Table 8), except cholesterol for which

concentrations were greater in 28-d-old calves born from dams fed fat supplements compared with those born from dams fed CON (3.29 v. 3.14 mmol/l; *P* = 0.02). Mean plasma concentrations of glucose, total protein, NEFA, insulin and TAG were not different between calves fed FC-0 or FC-2 (*P* > 0.10). Nevertheless, calves fed FC-2 had greater (*P* < 0.001) concentrations of albumin (28 d: 35.2 v. 28.9 g/l and 77 d: 39.1 v. 31.9 g/l) and cholesterol (28 d: 3.30 v. 3.17 mmol/l and 77 d: 2.26 v. 2.12 mmol/l) compared with those fed FC-0. Plasma urea N at 28 and 77 d of age was lower (*P* < 0.001) for calves fed FC-2 compared with those fed FC-0. At the end of the study, alkaline phosphatase was greater (600 v. 537 units/l) (*P* < 0.001) for calves fed FC-2 compared with calves fed FC-0.

Rumen fermentation parameters

Concentrations of ruminal pH, ammonia and individual and total VFA were not affected by DD, CS or their interactions (Table 9).

Discussion

Effect of prepartum diets

Contrary to our hypothesis, the type of FA fed prepartum did not have an impact on calf performance. Nevertheless, the




Fig. 1. Effect of prepartum dam diets (a) and calf starter (CS) diets (b) on weekly body weights (kg) of Holstein dairy calves fed CS containing no fat supplement (FC-0) or supplemented with 2% fat Ca-salt (FC-2). Calves were born from dams fed diets supplemented with no fat (control; CON), Ca-salts of soyabean oil (CSO) or Ca-salts of fish oil (CFO) starting at 3 weeks before expected calving date. Standard errors are 0.76 and 0.62 for prepartum dam diets and CS diets, respectively. Contrast of FAT = (CSO + CFO) v. CON, SFO = CSO v. CFO and CS = FC-0 v. FC-2. Effects in model: FAT: $P=0.02$; CS: $P<0.01$; week: $P<0.001$; dam diets \times week: $P=0.03$; CS \times week: $P<0.001$. * Difference between treatments at a given week ($P \leq 0.05$). † Trend between treatments at a given week ($P \leq 0.10$). (a) (—□—), CON; (—○—), CSO; (---△---), CFO. (b) (—□—), FC-0; (—○—), FC-2.

supplementation of fat regardless of the type of FA (rich in linoleic acid (CSO) or EPA and DHA (CFO)) had an impact on calf immune status and performance.

As expected, plasma concentrations of total IgG at birth were very low (≤ 0.32 g/l) and in most cases undetectable. Only one calf in the present study failed to attain at least 10 g/l IgG after colostrum feeding, and it was born from a dam fed CON. Because calves were fed colostrum from their corresponding dams, calves born from dams fed fat supplements consumed more IgG as their dams produced colostrum with higher IgG contents compared with dams fed CON⁽¹¹⁾. The greater IgG intake of calves born from dams fed fat supplements may have affected the observed better AEA and greater plasma IgG of those calves compared with the ones born from dams fed CON. In contrast, in beef cattle, calves suckling mature cows supplemented with oilseeds enriched in C18 : 2n-6 have similar serum IgG concentrations compared with calves suckling cows fed no oilseeds⁽³⁵⁾. Our findings support those of others who reported that calves born from dams fed fat supplements (saturated or rich in n-6 FA) tended to have greater serum concentrations of total IgG and better AEA after colostrum feeding compared with calves born from dams fed no supplemental fat⁽¹⁰⁾. Factors that may influence AEA include (1) time after birth

when colostrum is fed, (2) amount of IgG fed, (3) rate of gastric emptying and (4) absorptive capacity of enterocytes for IgG⁽¹⁰⁾. In the present study, feeding time was constant and adequate. Quigley *et al.*⁽³⁶⁾ recommended the ingestion of 150–200 g of IgG in the first 24 h to reduce the risk of failure of passive transfer. In the present study, calves born from dams fed fat supplements received 28% more IgG compared with those born from dams fed CON. The present study did not aim to investigate the specific absorptive capacity of enterocytes but a potential benefit of prepartum FA feeding on changing FA profile of enterocytes' membrane and consequently their absorptive capacity should not be ruled out.

In the present study, fat supplementation during late gestation did not affect CS intake but calves born from dams fed fat compared with those born from CON-fed dams had a better ADG preweaning, and it was maintained across the 77 d of the study. Therefore, the better BW gain observed in calves born from dams fed fat supplements resulted in a trend for an improved FE. The improved ADG and FE observed in calves born from dams fed fat may be due to meeting the metabolic demands for elongated PUFA⁽¹⁷⁾ and may not be due to a better digestibility of nutrients, as DD did not impact apparent nutrient digestibility and VFA production. Although hip and wither heights at birth were similar among calves born from dams fed any prepartum diet, calves born from dams fed fat supplements were taller than those born from dams fed CON at weaning (56 d) and the end of the study (77 d). This delayed impact of an *in utero* FA exposure on growth (skeletal and BW gain) maybe reflects a better health and immune status (greater plasma IgG and lower RT during the preweaning period) of calves born from dams fed fat supplements. Although prepartum feeding did not alter birth weight, a potential epigenetic effect of DD (e.g. fat supplementation) on post-birth calf metabolism cannot be ruled out. Indeed, others have also reported this potential mechanism with strategic feeding of FA during uterine life in dairy^(37,38) and beef cattle^(39,40).

Previous studies have reported that plasma concentrations of cholesterol can be altered by fat and/or FA supplementation in humans⁽⁴¹⁾, rodents⁽⁴²⁾ and in dairy cows⁽⁴³⁾. In addition, we recently found that plasma cholesterol was greater in prepartum cows fed fat (CSO or CFO) compared with cows fed CON⁽¹¹⁾. In the present study, cholesterol concentrations were higher in 28-d-old calves born from dams fed fat supplements compared with those born from dams fed CON. In contrast, Garcia *et al.*⁽¹³⁾ reported that plasma concentrations of cholesterol of calves born from dams fed fat (SFA or EFA) or specific types of FA did not differ from that of calves born from dams fed no fat supplement. Researchers have reported that cholesterol is transferred across the placenta, reaches fetal circulation, and thus substantially contributes to the fetal cholesterol pool in animals and humans^(44,45). Similar transfer mechanism may occur in calves born from dams fed fat supplements, and it may carry over the first weeks of life.

Effect of preweaning diets

In the present study, adding fat supplements to the CS did not affect intake of starter diets. Hill *et al.*⁽²²⁾ reported that preweaning calves fed a starter diet containing tallow or soyabean oil had

Table 5. Effect of dietary treatments on structural growth of Holstein calves fed a calf starter (CS) containing no fat supplement (FC-0) or supplemented with 2% fat calcium-salt (FC-2) during experimental periods (Mean values and pooled standard errors)

Items*	Dam diet and calf starters						SEM	P†		
	CON		CSO		CFO			Dam diet		CS
	FC-0	FC-2	FC-0	FC-2	FC-0	FC-2		FAT	SFO	
Hip height (cm)										
Day 3	82.9	83.1	83.3	83.3	82.9	83.1	0.51	0.81	0.58	0.65
Day 56	91.3	93.2	92.2	94.3	92.1	94.4	0.62	0.05	0.97	<0.01
Day 77	96.8	97.8	97.3	98.8	97.5	98.7	0.41	0.03	0.92	<0.01
Hip width (cm)										
Day 3	16.3	16.6	16.2	17.1	16.6	17.2	0.68	0.55	0.72	0.26
Day 56	21.0	21.8	21.5	22.2	21.7	22.1	0.62	0.39	0.91	0.18
Day 77	25.2	25.6	25.3	26.1	25.2	26.1	0.73	0.59	0.98	0.22
Heart girth (cm)										
Day 3	78.7	79.2	79.6	78.9	79.5	79.7	0.61	0.36	0.55	0.96
Day 56	90.9	91.1	91.5	90.8	90.1	91.8	0.75	0.94	0.77	0.47
Day 77	98.4	98.7	99.1	98.3	98.4	99.3	0.65	0.68	0.76	0.76
Body length (cm)										
Day 3	42.0	42.3	42.5	42.6	42.2	42.4	0.76	0.67	0.72	0.73
Day 56	53.6	53.7	53.5	53.9	53.9	53.8	0.63	0.90	0.79	0.85
Day 77	60.3	61.0	60.3	60.7	60.9	60.2	0.67	0.76	0.95	0.77
Body barrel (cm)										
Day 3	83.5	84.5	84.8	85.2	84.9	84.7	0.94	0.26	0.84	0.58
Day 56	109.8	110.7	109.9	111.1	109.5	110.3	0.88	0.86	0.42	0.22
Day 77	123.0	123.6	122.9	123.4	122.6	122.7	0.98	0.50	0.48	0.45
Wither height (cm)										
Day 3	77.9	78.8	79.1	78.6	79.5	80.0	0.70	0.12	0.19	0.63
Day 56	83.8	85.8	84.6	86.8	84.8	86.7	0.53	0.03	0.90	<0.01
Day 77	87.8	90.2	89.2	90.70	89.1	91.0	0.54	0.03	0.86	<0.01

CON, no fat supplement; CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition.

* Calves were weaned at 56 d of age.

† P values for orthogonal contrasts: FAT = control v. CSO + CFO; SFO = CSO v. CFO; CS = FC-0 v. FC-2. Interactions of FAT or SFO with CS were not significant for any parameter reported in this table ($P \geq 0.13$).

Table 6. Total time devoted to perform different behaviours during 8 h of observation of Holstein calves fed a calf starter (CS) containing no fat supplement (FC-0) or supplemented with 2% fat calcium-salt (FC-2) in pre- and postweaning periods (Mean values and pooled standard errors)

Items*	Dam diet and calf starters						SEM	P†		
	CON		CSO		CFO			Dam diet		CS
	FC-0	FC-2	FC-0	FC-2	FC-0	FC-2		FAT	SFO	
Eating (min)										
Day 35	33.5	34.8	35.2	30.2	31.6	36.3	2.86	0.73	0.66	0.89
Day 70	65.2	64.4	59.7	59.7	63.2	59.4	3.11	0.14	0.62	0.55
Ruminating (min)										
Day 35	114.0	112.3	108.3	113.3	111.5	113.5	1.85	0.25	0.86	0.84
Day 70	70.4	72.6	72.1	74.7	69.3	74.7	2.56	0.98	0.15	0.36
Standing (min)										
Day 35	124.3	131.1	127.7	130.1	118.2	114.4	11.20	0.60	0.26	0.84
Day 70	99.85	101.2	106.0	109.6	103.8	108.5	8.15	0.33	0.97	0.50
Lying (min)										
Day 35	191.5	183.0	188.5	186.7	199.7	193.0	14.29	0.70	0.53	0.62
Day 70	226.7	222.3	222.4	219.6	225.0	220.2	15.00	0.83	0.91	0.74
Non-nutritive oral behaviour‡ (min)										
Day 35	16.6	18.7	20.2	19.6	19.8	17.0	2.38	0.47	0.53	0.80
Day 70	15.9	19.6	19.8	16.4	18.5	16.6	1.84	0.83	0.77	0.72

CON, no fat supplement; CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition.

* Starter was offered from 3 d of age. Preweaning: 3–56 d. Postweaning: 57–77 d old.

† P values for orthogonal contrasts: FAT = control v. CSO + CFO; SFO = CSO v. CFO; CS = FC-0 v. FC-2. Interactions of FAT or SFO with CS were not significant for any parameter reported in this table ($P \geq 0.13$).

‡ Tongue rolling, licking surfaces or eating wood shavings.

Table 7. Effect of dietary treatments on rectal temperature, respiratory rate, health scores, and percentage of days with poor attitude, fever, diarrhoea, and nasal discharge, faecal score and days with diarrhoea of Holstein calves fed a calf starter (CS) containing no fat supplement (FC-0) or supplemented with 2% fat calcium-salt (FC-2) during experimental periods (Mean values and pooled standard errors)

Items*	Dam diet and calf starters						SEM	P†		
	CON		CSO		CFO			Dam diet		CS
	FC-0	FC-2	FC-0	FC-2	FC-0	FC-2		FAT	SFO	
Rectal temperature (°C)										
Prewearing	39.22	39.18	39.10	38.92	39.08	38.82	0.09	<0.01	0.36	0.04
Postweaning	39.32	38.95	39.27	38.93	39.22	38.98	0.08	0.64	0.95	<0.01
Respiration rate (breaths/min)										
Prewearing	46.4	46.0	46.7	48.0	47.8	47.9	1.15	0.17	0.66	0.74
Postweaning	55.9	57.0	57.5	54.7	58.8	58.5	2.22	0.64	0.24	0.72
Attitude score‡	0.32	0.30	0.31	0.25	0.31	0.24	0.04	0.41	0.99	0.16
Nasal discharge‡	0.14	0.11	0.10	0.08	0.11	0.08	0.03	0.34	0.89	0.45
Days with diarrhoea	7.50	5.80	7.20	5.00	7.40	5.30	0.62	0.42	0.67	<0.001
Faecal score‡	0.82	0.62	0.77	0.53	0.78	0.55	0.070	0.30	0.87	<0.001
Days with (%)§										
Poor attitude	18.4	15.4	16.6	13.4	16.5	12.3	2.80	0.35	0.82	0.14
Diarrhoea	19.2	17.1	18.3	15.3	18.2	15.4	2.39	0.52	0.99	0.19
Nasal discharge	6.80	4.60	6.50	3.78	5.90	3.25	1.94	0.66	0.76	0.13
Fever (first 21 d)	8.45	6.27	8.16	5.45	7.58	4.93	2.07	0.64	0.78	0.15

CON, no fat supplement; CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition.

* Starter was offered from 3 d of age. Prewearing: 3–56 d. Postweaning: 57–77 d old.

† P values for orthogonal contrasts: FAT = control v. CSO + CFO; SFO = CSO v. CFO; CS = FC-0 v. FC-2. Interactions of FAT or SFO with CS were not significant for any parameter reported in this table ($P \geq 0.24$).

‡ Attitude score scale: 0 = responsive, 1 = non-active, 2 = depressed, and 3 = moribund. Faeces score scale: 0 = firm faeces, no diarrhoea; 1 = soft faeces, no diarrhoea; 2 = mild diarrhoea; 3 = watery, severe diarrhoea. Nasal score scale: 0 = normal serous discharge; 1 = small amount of unilateral cloudy discharge; 2 = bilateral cloudy or excessive mucus discharge; 3 = copious bilateral mucopurulent discharge.

§ Percentage of days with health issue over a 77-d period unless indicated. Poor attitude and nasal discharge if score > 0, fever if temperature $\geq 39.5^\circ\text{C}$ and diarrhoea if score > 1.

similar starter intake; however, postweaning, calves fed soyabean oil consumed less starter. These results are in accordance with Kadkhoday *et al.*⁽⁴⁶⁾ who found that feeding starter diets with different ratios of linoleic and α -linolenic acids did not affect starter intake of Holstein dairy calves. Therefore, the better BW gain observed in the present study, without changes in starter or milk intake, resulted in an improved FE of calves fed FC-2 compared with those fed FC-0. The improved ADG and FE may be due to meeting the metabolic demands for elongated PUFA in FC-2 CS-fed calves⁽¹⁷⁾. Similarly, Hill *et al.*⁽²¹⁾ reported that dairy calves fed starter diets supplemented with Ca-salts of flaxseed oil, rich in α -linolenic acid, had increased ADG and FE compared with non-supplemented calves.

Wither and hip heights at weaning and the end of the study were also greater for calves fed FC-2 compared with those fed FC-0. Recent studies reported that *n*-3 FA have a regulatory role in neonatal bone turnover in humans and different animal models^(47,48). Kadkhoday *et al.*⁽⁴⁶⁾ found that calves fed a lower ratio of *n*-6 to *n*-3 FA had greater hip height compared with calves fed no supplemental fat. In contrast, others have reported that pre-weaned calves fed *n*-6 FA had similar skeletal growth compared with those fed SFA⁽⁴⁹⁾.

Several authors have reported that moderate concentrations of FA have beneficial effects on health and immune system function of dairy calves^(17,50,51). The acute inflammatory response is protective; however, an unresolved inflammation could lead to amplified inflammation and chronic diseases. Currently, it is well-understood that an acute inflammation has a temporal peak and should be resolved to restore tissue function⁽⁵²⁾. Calves fed

FC-2 had lower RT and lower overall faecal scores, compared with calves fed FC-0, which resulted in those calves having a fewer number of days with diarrhoea (–2 d) compared with calves fed FC-0. These findings support the hypothesis that one or more FA provided with the CF-2 diet (e.g. oleic or linoleic acid) have an anti-inflammatory effect (lower RT) and speed up the resolution of inflammation (fewer days with diarrhoea). Others have also reported that days with abnormal faecal scores were reduced by the inclusion of different FA blends ((butyrate, coconut oil and flax oil) or (butyrate, medium-chain FA and linolenic acid)) in the MR^(51,53). Similarly, others have reported a reduction in faecal scores when feeding starters with flaxseed oil or FA blend (coconut oil and canola oil) with the MR^(20,46). These beneficial effects of health-related parameters observed in calves fed FC-2 may have elicited a positive impact on their growth rate and FE.

Dairy cattle responses to different feed ingredients and feeding strategies differ. Indeed, it has been reported that fat content can affect eating behaviour⁽⁵⁴⁾. In the present study, animal behaviour was similar across treatments, and all calves spent similar times eating, ruminating, standing, lying or non-nutritive oral behaviour during both pre- and postweaning periods. To the best of our knowledge, animal behaviour has not been measured in experiments where fat supplements were fed *in utero* and preweaning and early postweaning simultaneously. Nevertheless, Kadkhoday *et al.*⁽⁴⁶⁾ reported that adding palm fat powder or flaxseed oil or both to starter diets had no adverse effect on the feeding behaviours of Holstein calves during pre- and postweaning

Table 8. Effect of dietary treatments on blood metabolites of Holstein calves fed a calf starter (CS) containing no fat supplement (FC-0) or supplemented with 2% fat calcium-salt (FC-2) during experimental periods (Mean values and pooled standard errors)

Items*	Dam diet and calf starters						SEM	P†		
	CON		CSO		CFO			Dam diet		CS
	FC-0	FC-2	FC-0	FC-2	FC-0	FC-2		FAT	SFO	
Glucose (mmol/l)										
Day 28	5.86	6.12	6.08	6.22	6.32	6.13	0.21	0.27	0.71	0.69
Day 56	4.90	4.92	4.89	5.04	4.84	5.02	0.11	0.69	0.77	0.22
Day 77	4.56	4.68	4.63	4.75	4.61	4.77	0.10	0.45	0.97	0.11
Total protein (g/l)										
Day 28	60.0	59.8	60.5	63.5	59.5	62.6	2.3	0.45	0.72	0.28
Day 56	61.4	64.3	63.0	65.2	63.3	65.7	2.1	0.44	0.82	0.15
Day 77	64.3	65.1	66.3	68.9	66.0	67.5	2.4	0.27	0.74	0.41
Albumin (g/l)										
Day 28	29.5	34.8	28.4	36.4	28.8	34.5	1.7	0.91	0.66	<0.001
Day 56	33.2	34.7	34.5	37.3	33.6	35.4	1.6	0.37	0.39	0.15
Day 77	31.2	38.8	32.6	39.5	32.0	39.1	1.7	0.63	0.80	<0.001
Urea N (mmol/l)										
Day 28	6.89	6.25	6.82	6.18	6.85	6.18	0.19	0.73	0.89	<0.001
Day 56	6.43	6.21	6.39	6.14	6.32	6.18	0.15	0.68	0.96	0.13
Day 77	7.92	7.19	7.89	7.28	7.93	7.28	0.23	0.80	0.94	<0.001
NEFA (mmol/l)										
Day 28	0.212	0.235	0.213	0.226	0.220	0.231	0.013	0.90	0.62	0.16
Day 56	0.168	0.180	0.176	0.164	0.181	0.174	0.010	0.98	0.48	0.82
Day 77	0.173	0.177	0.183	0.182	0.179	0.182	0.009	0.38	0.81	0.76
Insulin (μIU/ml)										
Day 28	27.0	26.7	28.1	27.9	26.7	27.4	0.70	0.25	0.16	0.90
Day 56	24.9	25.1	25.4	26.4	25.1	26.3	0.78	0.26	0.77	0.20
Day 77	16.8	18.1	16.4	17.1	17.5	17.2	0.87	0.50	0.51	0.41
Cholesterol (mmol/l)										
Day 28	3.09	3.19	3.21	3.37	3.21	3.36	0.07	0.02	0.96	0.02
Day 56	2.94	2.98	2.94	3.01	2.91	3.00	0.05	0.76	0.70	0.29
Day 77	2.12	2.26	2.11	2.26	2.13	2.27	0.05	0.91	0.73	<0.001
TAG (mmol/l)										
Day 28	0.26	0.25	0.25	0.25	0.25	0.26	0.01	0.30	0.45	0.50
Day 56	0.25	0.24	0.24	0.24	0.24	0.24	0.01	0.26	0.90	0.62
Day 77	0.21	0.20	0.21	0.20	0.20	0.20	0.01	0.21	0.40	0.23
Alkaline phosphatase (units/l)										
Day 28	553	573	568	581	557	585	18.3	0.53	0.82	0.18
Day 56	567	575	565	571	547	580	17.3	0.71	0.78	0.27
Day 77	531	598	540	600	542	603	19.6	0.69	0.94	<0.001

CON, no fat supplement; CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition.

* Calves were weaned at 56 d of age.

† P values for orthogonal contrasts: FAT = control v. CSO + CFO; SFO = CSO v. CFO; CS = FC-0 v. FC-2. Interactions of FAT or SFO with CS were not significant for any parameter reported in this table ($P \geq 0.22$).

periods. Furthermore, Benson *et al.*⁽⁵⁵⁾ reported that dairy cows infused, via the abomasum, with a mixture of sunflower and rapeseed oils did not alter their daily time spent engaged in eating, ruminating and idling activities. Similarly, McDonnell *et al.*⁽⁵⁶⁾ found that feeding behaviour of bull calves was not altered with fish oil supplementation during the preweaning period.

Studies have reported that strategic feeding of fat and FA alter the plasma metabolic profile of supplemented animals^(13,17) and the present findings are in support of those studies. Indeed, the feeding of a CS rich in unsaturated FA (FC-2) compared with a starter without supplemental fat (FC-0) altered the plasma metabolic profile of calves. The liver plays a key role in nutrient metabolism and utilisation due to its strategic position of integration within the circulatory system⁽³⁸⁾. Garcia *et al.*⁽³⁸⁾ reported that calves fed diets rich in *n*-6 compared with SFA up-regulated

the expression of hepatic genes involved in protein synthesis. In the present study, calves fed FC-2, compared with those fed FC-0, had higher plasma albumin concentrations, which may suggest a better hepatic function. Similarly, the addition of an FA blend to MR showed an increase in serum albumin concentration⁽⁵³⁾. The blood urea N-reducing effect of feeding FC-2 compared with FC-0 may be due to an improvement in protein deposition and reduction in protein oxidation, which is supported by higher concentrations of albumin and better ADG and skeletal growth of calves fed FC-2. Moreover, Le Roith *et al.*⁽⁵⁷⁾ reported that lower urea N could be related to changes in protein synthesis and protein oxidation, which are confirmed by our present findings.

Cholesterol is synthesised primarily in the liver of primuminant calves as products of lipid metabolism⁽⁵⁸⁾. The observed higher concentrations of cholesterol in plasma of calves fed

Table 9. Effect of dietary treatments on ruminal parameters of Holstein calves fed a calf starter (CS) containing no fat supplement (FC-0) or supplemented with 2% fat calcium-salt (FC-2) during experimental periods (Mean values and pooled standard errors)

Items*	Dam diet and calf starters						SEM	P†		
	CON		CSO		CFO			Dam diet		CS
	FC-0	FC-2	FC-0	FC-2	FC-0	FC-2		FAT	SFO	
pH										
Day 30	5.44	5.52	5.42	5.54	5.46	5.57	0.13	0.86	0.83	0.35
Day 70	5.65	5.67	5.61	5.73	5.64	5.74	0.11	0.78	0.85	0.39
NH ₃ -N (mg/dl)										
Day 30	11.1	9.80	11.4	12.0	11.5	10.8	1.24	0.36	0.65	0.64
Day 70	12.9	13.9	14.4	15.0	14.5	15.7	1.32	0.19	0.76	0.38
Acetate (mmol/l)										
Day 30	47.0	49.0	48.7	49.9	46.5	48.3	1.65	0.80	0.24	0.19
Day 70	49.6	50.6	50.7	51.3	48.3	49.9	1.85	0.97	0.31	0.47
Propionate (mmol/l)										
Day 30	38.2	40.2	39.9	41.1	37.7	39.5	1.58	0.80	0.24	0.20
Day 70	41.5	42.5	42.6	43.2	40.2	42.0	1.83	0.99	0.33	0.44
Butyrate (mmol/l)										
Day 30	14.0	15.4	15.7	15.8	13.5	15.3	1.98	0.84	0.51	0.50
Day 70	16.8	17.8	17.9	17.1	15.5	17.3	2.03	0.84	0.59	0.67
Valerate (mmol/l)										
Day 30	2.70	2.20	2.92	3.84	3.01	3.60	0.73	0.16	0.92	0.57
Day 70	3.15	3.30	3.23	3.62	3.32	3.00	0.73	0.92	0.72	0.90
Isovalerate (mmol/l)										
Day 30	0.72	0.85	0.83	0.78	0.73	0.60	0.15	0.72	0.33	0.92
Day 70	0.66	0.85	0.90	0.72	0.74	1.02	0.18	0.55	0.63	0.48
Total VFA (mmol/l)										
Day 30	102.7	107.8	108.1	111.5	101.5	107.5	5.07	0.66	0.29	0.25
Day 70	115.6	123.0	119.0	121.0	112.0	118.2	8.18	0.75	0.54	0.51
A:P										
Day 30	1.30	1.22	1.23	1.22	1.19	1.25	0.075	0.60	0.92	0.85
Day 70	1.17	1.15	1.20	1.16	1.18	1.15	0.062	0.69	0.79	0.60

CON, no fat supplement; CSO, supplementation with 140 g/d Ca-salts of soyabean oil; CFO, supplementation with 140 g/d Ca-salts of fish oil during the last 3 weeks before expected parturition; VFA, volatile fatty acids; A:P, acetate:propionate ratio.

* Calves were weaned at 56 d of age.

† P values for orthogonal contrasts: FAT = control v. CSO + CFO; SFO = CSO v. CFO; CS = FC-0 v. FC-2. Interactions of FAT or SFO with CS were not significant for any parameter reported in this table ($P \geq 0.21$).

FC-2 may simply reflect the preferential esterification of cholesterol esters with high intake of unsaturated FA compared with that of calves fed FC-0. Previous studies in preweaned calves reported different impacts of FA supplemented with MR, i.e. a linoleic acid-enriched MR reduced plasma cholesterol concentrations compared with a lauric acid-enriched MR⁽¹³⁾, whereas a soyabean oil-enriched MR quadratically increased plasma concentrations of cholesterol⁽¹⁷⁾.

Hill *et al.*^(59,60) reported that the increase in ALP is a good index of bone formation in young calves. Also, other researchers reported that adding flaxseed oil in the starter diet increased ALP^(21,44). In our study, plasma concentrations of ALP were greater in 77-d-old calves fed FC-2, compared with those fed CF-0. The use of ALP as an index of bone formation is supported by our findings of better skeletal growth (hip and wither heights) of calves fed FC-2 compared with those fed CF-0.

Several authors have reported that high amounts of PUFA in the rumen of mature cattle have adverse effects on micro-organism function and led to disturbed ruminal fermentation^(61,62). The lack of effect of CS on rumen fermentation parameters observed in the present study supports that of Kadkhoday *et al.*⁽⁴⁶⁾ who reported that CS supplemented with low levels of palm or flaxseed oil (less than 3% starter diet) had no adverse effect on ruminal fermentative parameters. On the other hand,

Ghasemi *et al.*⁽⁴⁹⁾ reported that feeding a starter diet with different types of fats had minor impacts on rumen fermentation parameters, e.g. ruminal pH, ammonia, or total VFA concentration were not altered. However, rumen fluid from calves fed a mixture of fats (palm fat, soybean oil and fish oil), compared with calves fed no fat or a single fat source, had increased molar concentrations of propionate and lower acetate:propionate ratio in ruminal fluid of 35-d-old Holstein calves⁽⁴⁹⁾. Furthermore, preweaned dairy calves fed a low-fat starter (2.9%) tended to have a reduction in the molar proportion of rumen butyrate but not in the other VFA⁽⁶³⁾. The mean ruminal pH values were within normal physiological values and ranged from 5.4 to 6.2, reflecting no adverse effect of diets on ruminal fermentation. Others also reported that extruded full-fat soyabeans did not alter ruminal pH in dairy calves⁽⁶⁴⁾. In the present study, ether extract content in CF-2 (rich in palmitic, oleic and linoleic acids) was 5.1% compared with 3.35% in CF-0 and these differences did not alter ammonia, specific VFA, or total VFA concentrations, which may suggest that cellulolytic bacterial populations were not affected by the difference in ether extract content and FA profile.

Finally, as consumer awareness for nutraceutical food has risen, nutritionists should focus on strategic feeding aiming to produce food from animal origin rich in nutrients conferring health benefits. Indeed, *n*-3 and *n*-6 FA are well known to have an

impact on human health and incidence of chronic diseases. Therefore, feeding PUFA to dairy cattle will not only benefit the performance of supplemented animals but their food products (e.g. milk, meat) would also have a FA profile with beneficial effects on human health.

Conclusions

Strategic feeding of diets enriched in PUFA (*n*-6 or *n*-3 FA) to pregnant non-lactating dams (1.15% of dietary DM) or a starter rich in unsaturated FA to preweaned Holstein calves (2% of CS DM) impacted calf performance. Prepartum supplementation of fat improved calves' passive immunity, overall ADG, weaning weight, final weight, hip height and wither height of calves without changing DMI. However, prepartum diets did not influence the effect of CS on calf performance or other measured parameters in the present study. Calves fed a CS enriched in unsaturated FA compared with non-supplemented calves had better ADG and FE, greater hip height, lower RT, fewer days with diarrhoea and higher plasma albumin and ALP. Further research is needed to increase the understanding of the effects of fat supplemented during the last weeks of uterine life and preweaning period on inflammatory status and response of immune cells of preweaned calves.

Acknowledgements

This research was supported by funds from the Agricultural Sciences and Natural Resources University of Khuzestan and University of Tehran. The authors thank the manager (Dr M. A. Moshayedi) and staff of the Zarin Khoshe dairy farm in the central part of Iran (Arak Province).

The authors' contributions were as follows: T. M., M. D., M. G. and M. C. designed the experiment; A. R. J. performed the research, laboratorial analyses and analysing the data. A. R. J. and M. G. wrote the manuscript. All authors reviewed and edited the manuscript.

The authors declare that there are no conflicts of interest.

References

- Soberon F & Van Amburgh M (2013) Lactation Biology Symposium: the effect of nutrient intake from milk or milk replacer of preweaned dairy calves on lactation milk yield as adults: a meta-analysis of current data. *J Anim Sci* **91**, 706–712.
- Gelsinger S, Heinrichs A & Jones C (2016) A meta-analysis of the effects of preweaned calf nutrition and growth on first-lactation performance. *J Dairy Sci* **99**, 6206–6214.
- Burr GO & Burr MM (1930) On the nature and role of the fatty acids essential in nutrition. *J Biol Chem* **86**, 587–621.
- Duttaroy AK (2000) Transport mechanisms for long-chain polyunsaturated fatty acids in the human placenta. *Am J Clin Nutr* **71**, 315S–322S.
- Neuringer M, Anderson GJ & Connor WE (1988) The essentiality of *n*-3 fatty acids for the development and function of the retina and brain. *Annu Rev Nutr* **8**, 517–541.
- Anderson GJ & Connor WE (1989) On the demonstration of omega-3 essential-fatty-acid deficiency in humans. *Am J Clin Nutr* **49**, 585–587.
- Simons K & Ikonen E (1997) Functional rafts in cell membranes. *Nature* **387**, 569–572.
- Clarke SD (2000) Polyunsaturated fatty acid regulation of gene transcription: a mechanism to improve energy balance and insulin resistance. *Br J Nutr* **83**, 59–866.
- Sampath H & Ntambi JM (2005) Polyunsaturated fatty acid regulation of genes of lipid metabolism. *Annu Rev Nutr* **25**, 317–340.
- Garcia M, Greco LF, Favoreto MG, *et al.* (2014) Effect of supplementing fat to pregnant nonlactating cows on colostral fatty acid profile and passive immunity of the newborn calf. *J Dairy Sci* **97**, 392–405.
- Jolazadeh AR, Mohammadabadi T, Dehghan-banadaky M, *et al.* (2019) Effect of supplementing calcium salts of *n*-3 and *n*-6 fatty acid to pregnant nonlactating cows on colostrum composition, milk yield, and reproductive performance of dairy cows. *Anim Feed Sci Technol* **247**, 127–140.
- Moallem U & Zachut M (2012) Short communication: the effects of supplementation of various *n*-3 fatty acids to late-pregnant dairy cows on plasma fatty acid composition of the newborn calves. *J Dairy Sci* **95**, 4055–4058.
- Garcia M, Greco LF, Favoreto MG, *et al.* (2014) Effect of supplementing essential fatty acids to pregnant nonlactating Holstein cows and their preweaned calves on calf performance, immune response, and health. *J Dairy Sci* **97**, 5045–5064.
- Lake SL, Scholljegerdes EJ, Weston TR, *et al.* (2006) Postpartum supplemental fat, but not maternal body condition score at parturition, affects plasma and adipose tissue fatty acid profiles of suckling beef calves. *J Anim Sci* **84**, 1811–1819.
- Jenkins KJ, Kramer JK, Sauer G, *et al.* (1985) Influence of triglycerides and free fatty acids in milk replacers on calf performance, blood plasma, and adipose lipids. *J Dairy Sci* **68**, 669–680.
- Jenkins KJ & Kramer JKG (1986) Influence of low linoleic and linolenic acids in milk replacer on calf performance and lipids in blood plasma, heart, and liver. *J Dairy Sci* **69**, 1374–1386.
- Garcia M, Shin JH, Schlaefli A, *et al.* (2015) Increasing intake of essential fatty acids from milk replacer benefits performance, immune responses, and health of preweaned Holstein calves. *J Dairy Sci* **98**, 458–477.
- Salehi R, Ambrose DJ & Oba M (2016) Short communication: effects of prepartum diets supplemented with rolled oilseeds on Brix values and fatty acid profile of colostrum. *J Dairy Sci* **99**, 3598–3601.
- Quigley JD III & Drewry JJ (1998) Nutrient and immunity transfer from cow to calf pre- and post-calving. *J Dairy Sci* **81**, 2779–2790.
- Hill T, Aldrich J, Schlotterbeck R, *et al.* (2007) Effects of changing the fat and fatty acid composition of milk replacers fed to neonatal calves. *Prof Anim Sci* **23**, 135–143.
- Hill T, Bateman H, Aldrich J, *et al.* (2009) Effects of changing the essential and functional fatty acid intake of dairy calves. *J Dairy Sci* **92**, 670–676.
- Hill TM, Bateman HG, Aldrich JM, *et al.* (2015) Inclusion of tallow and soybean oil to calf starters fed to dairy calves from birth to four months of age on calf performance and digestion. *J Dairy Sci* **98**, 4882–4888.
- Federation of Animal Science Societies (2010) *Guide for the Care and Use of Agricultural Animals in Research and Teaching*, 3rd ed. Champaign, IL: Federation of Animal Science Societies.
- Tyler JW, Hancock DD, Parish SM, *et al.* (1996) Evaluation of 3 assays for failure of passive transfer in calves. *J Vet Intern Med* **10**, 304–307.
- Quigley JD, Drewry IIIJ & Martin KR (1998) Estimation of plasma volume in Holstein and Jersey calves. *J Dairy Sci* **81**, 1308–1312.

26. Lesmeister KE & Heinrichs AJ (2005) Effects of adding extra molasses to a texturized calf starter on rumen development, growth characteristics, and blood parameters in neonatal dairy calves. *J Dairy Sci* **88**, 411–418.
27. Association of Official Analytical Chemists (1990) *Official Methods of Analysis*, 15th ed. Washington, DC: AOAC.
28. Van Soest PJ, Robertson J & Lewis B (1991) Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci* **74**, 3583–3597.
29. Van-Keulen J & Young BA (1977) Evaluation of acid-insoluble ash as a natural marker in ruminant digestibility studies. *J Anim Sci* **44**, 282–289.
30. Crooke W & Simpson W (1971) Determination of ammonium in Kjeldahl digests of crops by an automated procedure. *J Sci Food Agric* **22**, 9–10.
31. Castells L, Bach A, Araujo G, *et al.* (2012) Effect of different forage sources on performance and feeding behavior of Holstein calves. *J Dairy Sci* **95**, 286–293.
32. Yari M, Nikkhab A, Alikhani M, *et al.* (2010) Physiological calf responses to increased chromium supply in summer. *J Dairy Sci* **93**, 4111–4120.
33. SAS (2003) *SAS User's Guide: Statistics*, version 9.0. Cary, NC: SAS Institute.
34. Kenward MG & Roger JH (1997) Small sample inference for fixed effects from the restricted maximum likelihood. *Biometrics* **53**, 983–997.
35. Dietz RE, Hall JB & Whittier WD, *et al.* (2003) Effects of feeding supplemental fat to beef cows on cold tolerance in newborn calves. *J Anim Sci* **81**, 885–894.
36. Quigley J, Kost CJ & Wolfe TM (2002) Absorption of protein and IgG in calves fed a colostrum supplement or replacer. *J Dairy Sci* **85**, 1243–1248.
37. Garcia M, Greco LF, Lock AL, *et al.* (2016) Programming effect of dietary fatty acids on performance of Holstein heifers from birth through first lactation. *Anim Feed Sci Technol* **222**, 64–74.
38. Garcia M, Greco LF, Lock AL, *et al.* (2016) Supplementation of essential fatty acids to Holstein calves during late uterine life and first month of life alters hepatic fatty acid profile and gene expression. *J Dairy Sci* **99**, 7085–7101.
39. Marques RS, Cooke RF, Rodrigues MC, *et al.* (2017) Effects of supplementing calcium salts of polyunsaturated fatty acids to late-gestating beef cows on performance and physiological responses of the offspring. *J Anim Sci* **95**, 5347–5357.
40. Funston RN, Larson DM & Vonnahme KA (2010) Effects of maternal nutrition on conceptus growth and offspring performance: implications for beef cattle production. *J Anim Sci* **88**, Suppl., E205–E215. 2009-2351.
41. Chan JK, Bruce VM & McDonald BE (1991) Dietary α -linolenic acid is as effective as oleic acid and linoleic acid in lowering blood cholesterol in normolipidemic men. *Am J Clin Nutr* **53**, 1230–1234.
42. Vijaimohan K, Jainu M, Sabitha KE, *et al.* (2006) Beneficial effects of alpha linolenic acid rich flaxseed oil on growth performance and hepatic cholesterol metabolism in high fat diet fed rats. *Life Sci* **79**, 448–454.
43. Drackley JK, Klusmeyer TH, Trusk AM, *et al.* (1992) Infusion of long-chain fatty acids varying in saturation and chain length into the abomasum of lactating dairy cows. *J Dairy Sci* **75**, 1517–1526.
44. Woollett LA (2005) Maternal cholesterol in fetal development: transport of cholesterol from the maternal to the fetal circulation. *Am J Clin Nutr* **82**, 1155–1161.
45. Yoshida S & Wada Y (2005) Transfer of maternal cholesterol to embryo and fetus in pregnant mice. *Lipid Res* **46**, 2168–2174.
46. Kadkhoday A, Riasi A, Alikhani M, *et al.* (2017) Effects of fat sources and dietary C18: 2 to C18: 3 fatty acids ratio on growth performance, ruminal fermentation and some blood components of Holstein calves. *Livest Sci* **204**, 71–77.
47. Lau BY, Cohen DJ, Ward WE, *et al.* (2013) Investigating the role of polyunsaturated fatty acids in bone development using animal models. *Molecules* **18**, 14203–14227.
48. Kajarabille N, Peña M, Díaz-Castro J, *et al.* (2018) Omega-3 LCPUFA supplementation improves neonatal and maternal bone turnover: a randomized controlled trial. *J Funct Foods* **46**, 167–174.
49. Ghasemi E, Azad-Shahraki M & Khorvash M (2017) Effect of different fat supplements on performance of dairy calves during cold season. *J Dairy Sci* **100**, 5319–5328.
50. Ballou MA, Cruz GD, Pittroff W, *et al.* (2008) Modifying the acute phase response of Jersey calves by supplementing milk replacer with omega-3 fatty acids from fish oil. *J Dairy Sci* **91**, 3478–3487.
51. Hill T, VandeHaar M, Sordillo L, *et al.* (2011) Fatty acid intake alters growth and immunity in milk-fed calves. *J Dairy Sci* **94**, 3936–3948.
52. Serhan CN & Levy BD (2018) Resolvins in inflammation: emergence of the pro-resolving superfamily of mediators. *J Clin Invest* **128**, 2657–2669.
53. Esselburn KM, O'Diam KM, Hill TM, *et al.* (2013) Intake of specific fatty acids and fat alters growth, health, and titers following vaccination in dairy calves. *J Dairy Sci* **96**, 5826–5835.
54. Albright JL (1993) Feeding behavior of dairy cattle. *J Dairy Sci* **76**, 485–498.
55. Benson JA, Reynolds CK, Humphries DJ, *et al.* (2001) Effects of abomasal infusion of long-chain fatty acids on intake, feeding behavior and milk production in dairy cows. *J Dairy Sci* **84**, 1182–1191.
56. McDonnell RP, O'Doherty JV, Earley B, *et al.* (2019) Effect of supplementation with *n*-3 polyunsaturated fatty acids and/or β -glucans on performance, feeding behaviour and immune status of Holstein Friesian bull calves during the pre- and post-weaning periods. *J Anim Sci Biotechnol* **10**, 7.
57. Le Roith T, Hammond S, Todd S, *et al.* (2011) A modified live PRRSV vaccine and the pathogenic parent strain induce regulatory T cells in pigs naturally infected with *Mycoplasma hyopneumoniae*. *Vet Immunol Immunopathol* **140**, 312–316.
58. Drackley JK (2005) Interorgan lipid and fatty acid metabolism in growing ruminants. In *Biology of Metabolism in Growing Animals*, pp. 323–350 [DG Burrin and HJ Mersmann, editors]. Edinburgh, UK: Elsevier Limited.
59. Hill T, Aldrich J, Schlotterbeck R, *et al.* (2007) Amino acids, fatty acids, and fat sources for calf milk replacers. *Prof Anim Sci* **23**, 401–408.
60. Hill T, Aldrich J, Schlotterbeck R, *et al.* (2007) Effects of changing the fatty acid composition of calf starters. *Prof Anim Sci* **23**, 665–671.
61. Liu S, Bu D, Wang J, *et al.* (2012) Effect of incremental levels of fish oil supplementation on specific bacterial populations in bovine ruminal fluid. *J Anim Physiol Anim Nutr* **96**, 9–16.
62. Ivan M, Petit H, Chiquette J, *et al.* (2013) Rumen fermentation and microbial population in lactating dairy cows receiving diets containing oilseeds rich in C-18 fatty acids. *Br J Nutr* **109**, 1211–1218.
63. Kazemi-Bonchenari M, Mirzaei M, Jahani-Moghadam M, *et al.* (2016) Interactions between levels of heat-treated soybean meal and prilled fat on growth, rumen fermentation, and blood metabolites of Holstein calves. *J Anim Sci* **94**, 4267–4275.
64. ZeidAli-Nejad A, Ghorbani GR, Kargar S, *et al.* (2018) Nutrient intake, rumen fermentation and growth performance of dairy calves fed extruded full-fat soybean as a replacement for soybean meal. *Animal* **12**, 733–740.